

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 25 May 2016		2. REPORT TYPE Briefing Charts		3. DATES COVERED (From - To) 22 April 2016 - 25 May 2016	
4. TITLE AND SUBTITLE Theoretical Studies of nanocluster formation				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Robert J. Buszek, Jerry A. Boatz				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER Q188	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQRP 10 E. Saturn Blvd. Edwards AFB, CA 93524-7680				8. PERFORMING ORGANIZATION REPORT NO.	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQR 5 Pollux Drive Edwards AFB, CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-RQ-ED-VG-2016-097	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited. The U.S. Government is joint author of the work and has the right to use, modify, reproduce, release, perform, display, or disclose the work.					
13. SUPPLEMENTARY NOTES For presentation at AFOSR Molecular Dynamics and Theoretical Chemistry Program Review; Arlington, VA (25 May 2016) PA Case Number: #16215; Clearance Date: 5/5/2016					
14. ABSTRACT Viewgraph/Briefing Charts					
15. SUBJECT TERMS N/A					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 17	19a. NAME OF RESPONSIBLE PERSON J. Boatz
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NO (include area code) N/A



Theoretical studies of nanocluster formation

26 May 2016

Jerry Boatz

Principal Research Chemist

Aerospace Systems Directorate, RQRP

Air Force Research Laboratory

This briefing contains information up to:

Distribution A: Approved for public release; Distribution Unlimited. PA Clearance No: 16215



Outline



1. Introduction

- background, technical approach

2. Core-shell nanoclusters (Mg/Cu, Si/Al, etc.)

- energetic additives for propellants, explosives
- gas generators
- biocidal defeat agents

3. Summary and conclusions




Core-shell nanocluster synthesis



Core-shell nanoclusters such as SiAl_n , Ni_nAl_m , $\text{Al}_n(\text{CuO})_m$, etc. may be useful ingredients in propellants and explosives

- higher energy densities than organics ($\sim 3x$ RDX)
- some are resistant to surface oxidation (i.e., “magic clusters”)



Helium droplet
experiments at
AFRL/RW

Can core-shell nanoclusters be formed under cryogenic conditions (i.e., in helium droplet experiments) via stepwise condensation; i.e., what are the energy barriers (if any) to stepwise addition of atomic Al?






Cu_xMg_y core-shell nanocluster inversion



In helium droplet experiments, Mg atoms were captured in first pickup cell, followed by capture of Cu atoms to form Cu_xMg_y core-shell nanoclusters. However, scanning transmission electron microscopy (STEM) measures show cluster inversion occurred to produce Mg_yCu_x (!)




a) copper atoms

b) magnesium atoms

c) oxygen atoms

d) composite image






Cu_2Mg_{30}



1. Structure of Mg_{30} cluster was fully optimized.
2. Two Cu atoms were placed on opposite sides of Mg_{30} and structure reoptimized.
3. Distance between Cu atoms was decreased in steps of 0.25 Å, held fixed, and remaining DOF reoptimized.
4. Total energy plotted as function of fixed Cu-Cu distance.





DFT calculations: B3PW91/aug-cc-pwCVTZ(-PP) level



Cu_2Mg_{30} vs. Mg_{30}

Drop in energy in Cu_2Mg_{30} is not due to Mg_{30} rearranging to more stable structure.








Mg_2Cu_{30} vs. Cu_{30}

Drop in energy in Mg_2Cu_{30} is not due to Cu_{30} rearranging to more stable structure.







Formation of SiAl_2

MRMP2(10e,12o)/aug-cc-pvtz level of theory


Cyclic isomers




5A_1 , -0.63 eV




3B_1 , -1.59 eV




5A_2 , -1.55 eV



$^3A''$, -1.80 eV



1A_1 , -2.00 eV




3B_2 , -2.04 eV




Formation of SiAl_2

MRMP2(10e,12o)/aug-cc-pvtz level of theory


Linear isomers




$^5\Pi_u$, -0.58 eV




$^1\Pi$, -0.64 eV




$^1\Sigma_g^+$, -0.89 eV



$^5\Pi$, -0.70 eV



$^1\Pi_u$, -1.28 eV



$^3\Pi$, -0.94 eV



Summary and Conclusions



- **Mg/Cu core-shell nanoclusters**
 - Helium droplet experiments show inversion of Cu_xMg_y clusters to Mg_yCu_x .
 - Cu atoms diffusing into Mg_{30} , and vice-versa, have been modeled using DFT.
 - Estimated barrier for Cu atoms to migrate into Mg_n is < 1 kcal/mol.
 - Estimated barrier for Mg atoms to migrate into Cu_n is 6 kcal/mol.
 - Calculations are consistent with observed Cu/Mg inversion.
- **SiAl_n clusters**
 - SiAl_2 has multiple local minima which are more stable than ground state SiAl ($^4\Sigma^-$) + Al (2P)
 - Cyclic (C_{2v} and C_s) and linear ($D_{\infty h}$ and $C_{\infty v}$)
 - Singlet, triplet, quintet states
 - At long SiAl -----Al separations, preferred approach is linear
 - Al-Si-Al: $^5\Pi_u$ can form without a barrier
 - Si-Al-Al: $^5\Pi$ can form without a barrier
 - Barriers for linear \leftrightarrow cyclic isomerizations TBD
 - In helium droplet environment, linear quintet states may be formed.



Acknowledgements

Core-shell nanoparticles

- Dr. Robert Buszek (ERC, Inc.)
- Dr. Sam Emery(NSWC-IH), Y. Xin (Florida State Univ.),
C.J. Ridge, B.K. Little, C.M. Lindsay (AFRL/RW)
J.M. Boyle (Dublin School, Dublin, New Hampshire)

AFOSR

Dr. Mike Berman


DoD HPCMP



Backup Slides



Mg_n benchmark calculations



Mg_n clusters

“Closed shell” atomic configuration $[(1s)^2(2s)^2(2p)^6(3s)^2]$ suggests that weak dispersion interactions will be important. Need to consider

- core-core and core-valence correlation
- correlation method (MP2, CC, DFT)
 - “active” electrons to be correlated in MP2, CC
- suitable DFT functional for larger Mg_n clusters (up to $n \approx 100$)

Method	cc-pwCVDZ	cc-pwCVTZ	cc-pwCVQZ
MP2	23.1 / 3.042	28.5 / 3.013	29.1 / 3.011
CCSD(T)	16.7 / 3.100	tbd / 3.064	tbd / 3.065
DFT/B3PW91	26.5 / 3.092	26.5 / 3.091	26.4 / 3.092
DFT/PBE	34.6 / 3.070	34.5 / 3.070	34.3 / 3.070
DFT/PBE0	31.5 / 3.078	31.5 / 3.078	31.4 / 3.078
DFT/M06	30.8 / 3.028	30.4 / 3.025	TBD / TBD
DFT/M11	19.2 / 3.134	TBD / TBD	TBD / TBD

Calculated binding energies used to determine size of helium droplet needed for evaporative cooling




$SiAl_n$




Calculate potential energy surfaces of stepwise atomic addition reactions;
e.g., $SiAl_n + Al \rightarrow SiAl_{n+1}$

Diatomic: $Si (^3P) + Al (^2P) \rightarrow SiAl (^4\Sigma^-)$


Triatomic: $SiAl (^4\Sigma^-) + Al (^2P) \rightarrow SiAl_2$ (triplet or quintet)



$^3\Sigma^-$, -1.28 eV




$^1\Sigma_g^+$, -1.70 eV




$^3\Sigma_g^-$, -1.93 eV


ZAPT(2)/aug-cc-pvtz level
of theory




5A_2 , -1.77 eV





1A_1 , -2.65 eV





3B_2 , -2.67 eV



$\text{SiAl} (^4\Sigma) + \text{Al} (^2P)$



$\text{AlSi} (^4\Sigma) + \text{Al} (^2P)$



MRMP2(4e,5o)/aug-cc-pVTZ